



Case report

[Translated article] Abstinence syndrome following ingestion of poppy seed tea: A case report

Síndrome de abstinencia por consumo de té de semillas de adormidera: caso clínico

Teresa Rovira Medina^{a,c,*}, Pablo Yanes Sánchez^{a,c}, Miriam Bullich Ramon^{b,c,d}, Maria Oliver Cervelló^{a,c} and Mònica Gómez-Valent^{a,c,e}

^a Servicio de Farmacia, Parc Taulí Hospital Universitari, Sabadell, Barcelona, España

^b Servicio de Neumología, Parc Taulí Hospital Universitari, Sabadell, Barcelona, España

^c Institut d'Investigació i Innovació Parc Taulí (I3PT-CERCA), Parc Taulí Hospital Universitari, Barcelona, España

^d Facultad de Farmacia y Ciencias de la Alimentación, Universitat Autònoma de Barcelona, Barcelona, España

^e Facultad de Medicina, Unidad Docente Parc Taulí, Universitat Autònoma de Barcelona, Barcelona, España

Introduction

Processed poppy seeds (*Papaver somniferum* L.) are used for culinary purposes worldwide. However, unprocessed poppy seeds may contain varying concentrations of morphine, codeine, and thebaine, and be used to prepare beverages with psychoactive properties.^{1,2} The literature describes cases of dependence and toxicity associated with the consumption of poppy seed infusions.^{2–5}

We present the management of a patient with pulmonary tuberculosis who had an incidental finding of withdrawal syndrome, highlighting the importance of reconciling medication, botanicals, and foods.

Case description

A 32-year-old Indian man living in Spain, with a language barrier, no known medical history, and no habitual medication. Of note, the patient reported consuming 5–6 units of alcohol per day and denied the use of other drugs (no toxicity test was performed). The patient had a 2-week history of cough with hemoptotic sputum, progressive dyspnea, bilateral pleuritic chest pain, and shivering, accompanied by progressive deterioration of general health and associated constitutional symptoms (1.70 m, 48 kg), with the remainder of the examination being normal.

On arrival, he was hemodynamically stable, with the following findings (Table 1): physical examination revealed bilateral rhonchi without signs of increased respiratory effort and with painful hepatomegaly. Viral PCR and antigenuria tests were negative. Blood (subsequently negative) and sputum cultures were obtained. Antibiotic coverage was initiated with meropenem (1 g/8 h) and amikacin (1000 mg in a single dose). Subsequently, the sputum tested positive on the Ziehl-Neelsen

test, so treatment was started with isoniazid (250 mg/24 h), rifampicin (600 mg/24 h), pyrazinamide (1000 mg/24 h), and intravenous ethambutol (800 mg/24 h).

The clinical picture was acute respiratory failure due to likely advanced-stage tuberculous bronchopneumonia, with extrapulmonary involvement (later ruled out by CT scan), and suspected respiratory superinfection.

After one day of admission, the patient developed tachycardia (HR >150) and tachypnea (RR 27), with sustained blood pressure and fever, despite the improvement in acute phase reactants. He also had profuse sweating, disorientation, piloerection, shortness of breath, bilious vomiting, and diarrhea. He was admitted to the intermediate respiratory care unit, where high-flow oxygen therapy was initiated.

The patient's relatives were interviewed and reported that he had no toxic habits, did not take any regular medication, and only consumed daily infusions. Thanks to the photographs provided, it was determined that the infusions were made from poppy seeds. A literature search in Pubmed (using the terms “poppy tea” and “poppy seed”) identified articles^{1–5} supporting the possibility of withdrawal symptoms.

Withdrawal symptoms usually begin at 6–12 h and peak at 36–72 h.⁶ However, the use of an enzyme inducer such as rifampicin would explain the rapid evolution of the picture.

In view of these findings, the impossibility of using methadone (due to the lack of an oral route), and the marked interaction with rifampicin, subcutaneous morphine (2.5 mg/12 h) was initiated. After 12 h, the patient showed clinical improvement, with resolution of tachycardia (HR 115), no signs of increased respiratory effort, and normal breathing (RR 16).

We decided to continue treatment with the same dose of morphine until discharge, given the difficulty of follow-up by the psychiatric service for subsequent withdrawal management with methadone, the interaction with antituberculosis treatment, and the good response to morphine.

DOI of original article: <https://doi.org/10.1016/j.farma.2024.11.003>.

* Corresponding author.

E-mail address: terovira19@gmail.com (T. Rovira Medina).

<https://doi.org/10.1016/j.farma.2025.01.005>

1130-6343/© 2024 Sociedad Española de Farmacia Hospitalaria (S.E.F.H.). Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1

Abnormal clinical, blood, and radiological parameters on arrival.

Parameter or test	Value or result
Temperature	39 °C
HR	140 bpm
SaO ₂	94%
RR	25 breaths/min
CRP	21.6 mg/dL (0–0.5)
Procalcitonin	12.4 ng/mL (0–0.5)
Aspartate aminotransferase	461 U/L (0–38)
Alanine aminotransferase	210 U/L (10–41)
Alkaline phosphatase	136 U/L (40–129)
Gamma-glutamyl transpeptidase	76 U/L (8–61)
Conjugated bilirubin	0.7 mg/dL (0.1–0.25)
Blood gas	Compensated respiratory alkalosis
Chest X-ray	Bilateral pulmonary infiltrates

HR, heart rate; RR, respiratory rate; CRP, C-reactive protein; SaO₂, baseline oxygen saturation; bpm, beats per minute.

Discussion

In Spain, the sale of opium poppies to the public is prohibited due to their toxicity.⁷ Schedule 1 narcotics include morphine, which is found in opium (concentrated poppy juice) and poppy straw, which includes all parts of the plant after harvest, except the seeds.^{8,9} The seeds are therefore excluded from this classification and are offered for sale, for example, on various websites. Various concentrations of morphine, codeine, and thebaine have been measured in unprocessed seed infusions, with one study reporting concentrations from seeds of <1–2.788 mg/kg, <1–247.6 mg, and <1–124 mg, respectively, depending on the supplier and extraction method.

The literature has described cases of poppy seed infusion abuse that required medical attention for detoxification.^{2–4} There have also been reports of death¹ and neonatal abstinence syndrome following maternal ingestion.⁵ In most cases, morphine levels were measured in the blood or urine of the patients, or in the infusion itself. The main treatment options used were buprenorphine, methadone, and oral morphine, with some patients requiring intravenous or intramuscular naloxone as emergency treatment.

However, in the case described, morphine was not measured in biological fluids, nor was it possible to place the patient in a detoxification circuit due to a lack of knowledge of his habits and his commitment to detoxification.

The Spanish Early Warning System (SEAT) is a national mechanism for the annual identification of new psychoactive substances and the dissemination of information on the consequences of their use and abuse. These substances include botanicals and extracts.¹⁰ The cases described suggest that it may be necessary to study poppy seed infusions as a possible source of narcotics and, consequently, implement the corresponding regulatory measures to control their production and sale.

Conclusion

Unlike other reported cases, the reason for this patient's admission was not intoxication or interest in detoxification, but rather an incidental finding, which was also a confounding element in that it could have been interpreted as a worsening of the patient's condition due to a lack of effectiveness or an adverse reaction to the established treatment.

This situation highlights the importance of a complete and early pharmacological anamnesis, including not only the usual medications but also controlled drugs, phytotherapy, and supplements.

Ethical responsibilities

All ethical responsibilities have been met, in particular the anonymization of the case. The authors obtained authorization or informed consent from all those involved in this case.

Funding

This study has not received specific funding from public sector agencies, commercial sector entities, or not-for-profit organizations.

Responsibilities and ceding of rights

All authors accept the responsibilities defined by the International Committee of Medical Journal Editors (available at <http://www.icmje.org/>).

In the event of publication, the authors grant exclusive rights of reproduction, distribution, translation, and public communication (by any means or medium, whether sound, audiovisual, or electronic) of our work to *Farmacia Hospitalaria* and, by extension, to the SEFH. For this purpose, a letter of ceding of rights will be signed at the time of submitting the paper through the online manuscript management system.

CRediT authorship contribution statement

Teresa Rovira Medina: Writing – review & editing, Writing – original draft, Validation, Supervision, Methodology, Investigation, Formal analysis, Conceptualization. **Pablo Yanes Sánchez:** Writing – review & editing, Writing – original draft, Validation, Supervision, Methodology, Formal analysis, Conceptualization. **Miriam Bullich Ramon:** Writing – review & editing, Writing – original draft, Validation, Supervision, Methodology, Formal analysis, Data curation. **Maria Oliver Cervelló:** Writing – review & editing, Writing – original draft, Validation, Methodology, Formal analysis. **Mònica Gómez Valent:** Writing – review & editing, Writing – original draft, Validation, Methodology.

Declaration of competing interest

None declared.

References

1. Powers D, Erickson S, Swortwood MJ. Quantification of morphine, codeine, and thebaine in home-brewed poppy seed tea by LC-MS/MS. *J Forensic Sci.* 2018;63(4): 1229–35. doi: [10.1111/1556-4029.13664](https://doi.org/10.1111/1556-4029.13664).
2. Spyres MB, van Wijk XMR, Lapoint J, Levine M. Two cases of severe opiate toxicity after ingestion of poppy seed tea. *Toxicol Commun.* 2018;2(1):102–4. doi: [10.1016/0006-291x\(75\)90482-9](https://doi.org/10.1016/0006-291x(75)90482-9).
3. Haber I, Pergolizzi Jr J, LeQuang JA. Poppy seed tea: a short review and case study. *Pain Ther.* 2019;8(1):151–5. doi: [10.1007/s40122-019-0113-5](https://doi.org/10.1007/s40122-019-0113-5).
4. Kauppila GR, Eagen KV. Opioid use disorder from poppy seed tea use: a case report. *Am J Med Case Rep.* 2023;26(24). doi: [10.12659/AJCR.938675](https://doi.org/10.12659/AJCR.938675) e938675.1–e938675.3.
5. Garcia MR, Swortwood MJ, Aune CN, Ahmad KA. Maternal poppy seed tea ingestion and ensuing neonatal abstinence syndrome. *Neonatology.* 2020;117(4):529–31. doi: [10.1159/000508446](https://doi.org/10.1159/000508446).
6. Garcia Villa A, Parra González A. Trastornos por Consumo de Alcohol y Otras Drogas. *Manual de Diagnóstico y Terapéutica Médica Hospital Universitario 12 de Octubre 9ª edición.* Madrid: MSD; 2023.
7. Orden SCO/190/2004. De 28 de enero, Por la que se Establece la Lista de Plantas cuya Venta al Público Queda Prohibida o Restringida por Razón de su Toxicidad. *Boletín Oficial del Estado*, n° 32, 6 de febrero de 2004. [consultado 12 May 2024]. Disponible en: <https://www.boe.es/boe/dias/2004/02/06/pdfs/A05061-05065.pdf>.
8. Convención Única de 1961 sobre Estupefacientes enmendada por el Protocolo de 1972 de Modificación de la Convención Única de 1961 sobre Estupefacientes. Naciones Unidas; 1966. Suscrita por España el día 27 de julio de 1961, [consultada

- 23 May 2023]. Disponible en: <https://profesionales.msd.es/manual-12-de-octubre/cardiovascular/>.
9. Lista de estupefacientes sometidos a fiscalización internacional. Lista Amarilla. Anexo a los Formularios A, B y C 58a edición. Junta Internacional de Fiscalización de Estupefacientes; agosto de 2019 [consultada 22 May 2024]. Disponible en: https://www.incb.org/documents/Narcotic-Drugs/Yellow_List/58th_Edition/Yellow_List-SPA.pdf.
10. Observatorio Español de las Drogas y las Adicciones. INFORME 2023. Alcohol, Tabaco y Drogas Ilegales en España, 2023. [consultado 12 May 2024]. Disponible en: <https://pnsd.sanidad.gob.es/>.